



Clinical Pharmacology & Toxicology Pearl of the Week

~Antihypertensives, Part 2: Cardio-selective Beta-blockers~

β-1 receptors:

- ✓ Predominantly located on cardiac myocytes, kidney cells, and adipocytes
- ✓ Stimulated by adrenergic signaling molecules: (epinephrine/norepinephrine or dopamine)
- ✓ Adrenergic stimulation of G-PCR → ↑cAMP → ↑ phosphorylated calcium channels → ↑ calcium release → sarcoplasmic reticulum release of calcium → muscle contraction/effector action
 - Cardiac myocytes: ↑ sinoatrial and AV node firing + ventricular muscle firing → increased contractility and heart rate
 - Kidney: ↑ smooth muscle release of **renin**
 - Adipocytes: ↑ lipolysis

β-2 receptors:

- ✓ Found on smooth muscles (GI tract, bronchi, detrusor muscles, uterus, seminal tract), pancreas (For insulin and glucagon secretion), eye
- ✓ Stimulation leads to:
 - Smooth muscle relaxation
 - Increased ocular pressures (increased aqueous humour production)
- ✓ Less relevance in hypertension management

β-1 receptor antagonists:

- Indications: Hypertension, coronary-artery disease, heart failure, arrhythmias, angina-pectoris
- Examples:
 - **Cardio-selective (β-1 selective):** Atenolol, bisoprolol, esmolol, metoprolol
 - **Non-selective:** Nadolol, propranolol, timolol
 - **Non-selective with alpha-1 antagonism:** Carvedilol, labetalol
- **Anti-hypertensive mechanism of action:** Decreased renin (Decreased RAAS activity) and decreased inotropy and chronotropy.
- Anti-anginal mechanism of action: As above; mostly through negative chronotropy and inotropy (Lower O₂ demand within myocardium)

Pharmacokinetics of cardio-selective β -blockers

	Absorption	Distribution	Metabolism	Elimination
Atenolol	50% T_{peak} : 2-4hr	V_d : 0.9-1.6L/Kg 6-16% protein bound	85-90% unaltered Small amounts of conjugation	85% renal $\frac{1}{2}$ life: 6-7hrs
Bisoprolol	>80% T_{peak} : 2-4hr	V_d : 2.9L/Kg 30% protein bound	CYP 3A4 – inactive metabolites	98% renal 50% unaltered, 50% metabolites $\frac{1}{2}$ life: 9-12hrs
Esmolol	IV only	V_d : Unknown	RBC esterases	~80% renal (<2% unaltered) $\frac{1}{2}$ life: 9 min
Metoprolol	~75% T_{peak} : 1-2hr	V_d : 4.2L/kg 11% protein bound	CYP 2D6 – inactive metabolites	>95% renal <5% unaltered $\frac{1}{2}$ life: 3-7hrs

- ✓ **Anti-hypertensive effect:** Flat response curve (increasing the dose will change chronotropic response, but does not cause further reduction of blood pressure)
- ✓ Prescribing situations: Hypertension and...
 - Less than 60 years old (signal towards stroke in people on β -blockers over the age of 60)
 - Coronary artery disease (Within the last 18 months)
 - Heart failure
 - Angina
 - Arrhythmia, including atrial fibrillation
- ✓ Combinations:
 - RAAS additive effect: β -blocker + ACEi/ARB, β -blocker + thiazide/thiazide-like diuretic
 - RAAS + vasodilation: β -blocker + CCB

Adverse effects/Toxicity:

- ✓ Side effects: bradycardia, hypotension, diarrhea, fatigue +/- depression
- ✓ Caution initiating/Escalating: Decompensated heart failure, pre-syncope, bradycardia
- ✓ Toxicity: bradycardia + hypotension, less β -1 selectivity in toxic ingestions leading to bronchospasm, hypoglycemia, altered mental status

References:

1. Wong, G.W., Boyda, H.N., Wright, J.M., 2016. Blood pressure lowering efficacy of beta-1 selective beta blockers for primary hypertension. Cochrane Database of Systematic Reviews. doi:10.1002/14651858.cd007451.pub2
2. Westfall TC, Westfall DP. Adrenergic Agonists and Antagonists. In: Brunton LL, Chabner BA, Knollmann BC. eds. Goodman & Gilman's: The Pharmacological Basis of Therapeutics, 12e. McGraw-Hill
3. Benowitz NL. Antihypertensive Agents. In: Katzung BG. eds. Basic & Clinical Pharmacology, 14e. McGraw-Hill;
4. Micromedex: <https://www-micromedexsolutions-com.ahs.idm.oclc.org/>
5. Drugbank: <https://go.drugbank.com/>

The Clinical Pharmacology (CP) physician consultation service is available Mon-Fri, 8am-5pm. The on-call physician is listed in ROCA on the AHS Insite page. CP consultations are also available through Netcare e-referral and Specialist Link. You can also find us in the [Alberta Referral Directory](#) (ARD) by searching "Pharmacology" from the ARD home page. Click [HERE](#) for more details about the service.

The Poison and Drug Information Service (PADIS) is available 24/7 for questions related to poisonings. Please call 1-800-332-1414 (AB and NWT) or 1-866-454-1212 (SK). Information about our outpatient Medical Toxicology Clinic can be found in [Alberta Referral Directory](#) (ARD) by searching "Toxicology" from the ARD home page.

More CPT Pearls of the Week can be found [HERE](#).

Created: February 26, 2021

Reviewed: March 10, 2025