



Clinical Pharmacology & Toxicology Pearl of the Week


~ Drug-induced QTc prolongation ~

What are the QT and QTc intervals?

- ✓ The QT interval is measured from an electrocardiogram (ECG) from the start of the Q wave to the end of the T wave or predicted intersection of the T wave downslope and the baseline.
- ✓ The QT interval corresponds to the time taken for ventricular depolarisation and repolarization.
- ✓ The QT interval changes with heart rate. Therefore, a corrected QT interval (QTc) is often used to allow for assessment of interval length independent of heart rate.
- ✓ QTc can be calculated by many formulas such as the Bazett, Fridericia, Framingham Linear Regression, and Rautaharju Formulae.
- ✓ Prolonged QTc is defined as a QTc >440ms in men or a QTc >460ms in women.

Why is QTc prolongation dangerous?

- ✓ QTc >500ms is an independent risk factor for developing Torsades des Pointes (TdP).
- ✓ For acute ingestions with QTc prolonging drugs, it has been suggested that the absolute QT interval (not QTc) is a better predictor of the risk of TdP.
- ✓ Values can be plotted on the QT interval nomogram (Figure 1, right) to evaluate for risk of TdP.



Mechanism of QTc prolongation on ECG:

- ✓ QTc prolongation occurs when phase 2 or 3 of the action potential is prolonged (Figure 2).
- ✓ Generally, slow potassium efflux due to potassium channel dysfunction prolongs the repolarization phases 2 and 3. This can be due to delayed rectifier potassium channel blockade, hypokalemia, and hypomagnesemia (causing intracellular hypokalemia).
- ✓ Additional mechanisms of prolonged QTc include hypocalcaemia slowing phase 2 of the action potential, as well as sodium channel blockade resulting in prolonged QRS (and resultant prolonged QTc).

Mechanism of drug induced QTc prolongation:

- ✓ In drug induced QTc prolongation, prolongation results due to blockade of the hERG delayed potassium rectifier channel.
- ✓ This channel plays a key role in phase 3 of the cardiac action potential, where repolarization occurs.
- ✓ The hERG delayed potassium rectifier channel contains a relatively large pore size with multiple hydrophobic pockets allowing for promiscuous binding of a wide range of xenobiotics within the channel. This accounts for the long list of medications implicated in prolonging the QTc interval.


Drugs causing QTc prolongation

- ✓ Many commonly used medications are associated with QTc prolongation including antiarrhythmics, antibiotics, antidepressants, anti-emetics, antihistamines, and antimalarials.
- ✓ A list of commonly used medications that cause QTc prolongation is included in Table 1.

Class of Drug	Examples
Antiarrhythmic	Amiodarone, Procainamide, Quinidine, Sotalol
Antipsychotics	Chlorpromazine, Clozapine, Haloperidol, Quetiapine, Risperidone
Antibiotics	Azithromycin, Clarithromycin, Erythromycin, Fluconazole, Ketoconazole, Ciprofloxacin, Levofloxacin, Moxifloxacin, Trimethoprim-Sulfamethoxazole
Antidepressants	Amitriptyline, Citalopram, Fluoxetine, Nortriptyline, Paroxetine, Sertraline, Venlafaxine
Anti-emetics	Ondansetron, Prochlorperazine

Table 1. Examples of commonly used medications associated with prolonged QTc. (Adapted from Ayad et al. 2010).⁶

Figure 2. Phases of the cardiac action potential and associated ion channels involved (Roden 2004)



✓ **Are all QTc prolonging drugs equally likely to cause Tdp?**

- ✓ Although many drugs are known to cause prolonged QTc, some drugs are more commonly reported to cause TdP. Examples can be found in Table 2 below.
- ✓ Visit crediblemeds.org for a searchable database of drugs for risk of QTc prolongation and TdP.

Class of Drug	Examples
Antiarrhythmics	Amiodarone, Procainamide, Quinidine, Sotalol
Antibiotics	Clarithromycin, Erythromycin
Antipsychotic agents	Chlorpromazine, Haloperidol,
Anti-emetics	Domperidone, Droperidol
Opioids	Methadone
Others	Arsenic trioxide, Cisapride

Treatment of a prolonged QTc:

- ✓ Identify and remove drugs associated with prolonged QTc from patient's profile.
- ✓ Avoid the use of other QTc prolonging medications.
- ✓ Identify and correct electrolyte abnormalities; particularly hypokalemia, hypocalcaemia and hypomagnesemia.
- ✓ Monitor for arrhythmias. If TdP develops, treat per ACLS guidelines. Consider magnesium sulphate empirically if serum magnesium concentration unknown. Consider continuous cardiac monitoring if QTc >500 ms. Consider serial ECGs every two to six hours until resolution.

References:

1. Isbister GK. Risk assessment of drug-induced QTC prolongation. Australian prescriber. 2015;38(1):20.
2. Nachimuthu S, Assar MD, Schussler JM. Drug-induced QTC interval prolongation: mechanisms and clinical management. Ther Adv Drug Saf. 2012 Oct; 3(5): 241-253.
3. Vandenberg JL, Perozo E & Allen TW. Towards a structural view of drug binding to hERG K⁺ channels. Trends Pharmacol Sci. 2017; 38:899-907.
4. Roden D. Drug-Induced Prolongation of the QTC Interval. N Engl J Med. 2004; 350:1013-1022.
5. Ayad RF, Assas MD, Simpson L, Garner JB, Schussler JM. Causes and management of drug-induced long QTC syndrome. Proc (Bayl Univ Med Cent). 2010 Jul; 23(3):250-255.

The Clinical Pharmacology (CP) physician consultation service is available Mon-Fri, 8am-5pm. The on-call physician is listed in ROCA on the AHS Insite page. CP consultations are also available through Netcare e-referral and Specialist Link. You can also find us in the [Alberta Referral Directory](#) (ARD) by searching "Pharmacology" from the ARD home page. Click [HERE](#) for more details about the service.

The Poison and Drug Information Service (PADIS) is available 24/7 for questions related to poisonings. Please call 1-800-332-1414 (AB and NWT) or 1-866-454-1212 (SK). Information about our outpatient Medical Toxicology Clinic can be found in [Alberta Referral Directory](#) (ARD) by searching "Toxicology" from the ARD home page.

More CPT Pearls of the Week can be found [HERE](#).