

## **COPD Medications**



	I Take:	My Dose	What it Does	Side Effects <sup>+</sup>	Comments
Reliever as needed	Airomir® Bricanyl® Salbutamol <b>Ventolin®</b>		<ul> <li>Relaxes muscles in the airways</li> <li>Works within minutes when needed</li> <li>Used before activity to prevent attacks</li> </ul>	<ul> <li>tremor(hands shake)</li> <li>fast heart rate</li> <li>nervousness</li> <li>headache</li> <li>weakness/dizziness</li> <li>sweating</li> </ul>	<ul> <li>Keep with you at all times for use</li> <li>During lung attacks/flare-ups</li> <li>Episodes of shortness of breath</li> <li>Episodes of difficulty breathing</li> <li>Use spacer with all MDIs*</li> </ul>
)r	Tudorza®  Seebri®  Spiriva®  Incruse®		Relaxes muscles in the airways	<ul><li> dry mouth/throat</li><li> constipation</li><li> trouble urinating</li><li> headache</li></ul>	<ul> <li>Avoid getting the powder in your eyes</li> <li>Avoid getting Respimat mist in your eyes</li> <li>If trouble urinating, stop medication and see your Doctor</li> </ul>
g Bronchodilator	Serevent® Onbrez® Oxeze®		Relaxes muscles in the airways	<ul><li>headache</li><li>tremor(hands shake)</li><li>fast heart rate</li><li>muscle cramps</li></ul>	Varied onset
Long acting	Anoro® Duaklir® Inspiolto® Ultibro®		Relaxes muscles in the airways	<ul> <li>dry mouth/throat</li> <li>trouble urinating</li> <li>headache</li> <li>tremor (hands shake)</li> <li>fast heart rate</li> <li>muscle cramps</li> </ul>	<ul> <li>2 medications in one device</li> <li>Once daily usage</li> <li>Avoid getting the powder/Mist in your eyes</li> </ul>

		I Take:	My Dose	What it Does	Side Effects+	Comments
Combination	Combination preventer	Advair® Breo® Symbicort®		<ul> <li>Regular use decreases inflammation (swelling) of the airways</li> <li>Relaxes muscles in the airways</li> <li>Meant only for people who experience frequent flare-ups</li> </ul>	<ul> <li>hoarse voice</li> <li>sore throat</li> <li>oral thrush</li> <li>headache</li> <li>tremor (hands shake)</li> <li>fast heart rate</li> <li>muscle cramps dry mouth/throat</li> <li>trouble urinating</li> </ul>	<ul> <li>Rinse, gargle, and spit after each use</li> <li>Take regularly to control inflammation and relax muscles in airway</li> <li>Carries a slight increased risk of pneumonia</li> <li>Use spacer with all MDIs*</li> </ul>
	preventer + bronchodialator	Trelegy®		<ul> <li>Regular use decreases inflammation (swelling) of the airways</li> <li>Relaxes muscles in the airways</li> <li>Long acting once/day medication (24hrs)</li> </ul>	<ul> <li>hoarse voice</li> <li>sore throat</li> <li>oral thrush</li> <li>headache</li> <li>tremor (hands shake)</li> <li>fast heart rate</li> <li>muscle cramps</li> </ul>	<ul> <li>Rinse, gargle, and spit after each use</li> <li>Avoid getting medication into eyes</li> <li>Take regularly to control inflammation and relax muscles in airway</li> <li>Carries a slight increased risk of pneumonia</li> <li>If trouble urinating, stop use and see your Doctor</li> </ul>
10 d O	Snort-acting bronchodilator	Atrovent® Combivent®		<ul> <li>Relaxes muscles in the airways</li> <li>Starts working within 30 minutes</li> <li>Lasts 6 hours</li> </ul>	<ul> <li>dry mouth, bad taste</li> <li>tremor (hands shake)</li> <li>headache</li> <li>blurred vision</li> <li>stuffy nose</li> <li>trouble urinating</li> </ul>	<ul> <li>Rinse mouth with water if bothered by dryness or bad taste in mouth</li> <li>Avoid getting medication into eyes</li> <li>If trouble urinating, stop use and see your Doctor</li> <li>Use spacer with all MDIs*</li> <li>Combivent contains Atrovent and Ventolin</li> </ul>
Potent Anti-	inflammatory	Prednisone		<ul> <li>Rapidly decreases inflammation of the airways</li> <li>Usually taken for 5-10 days</li> </ul>	<ul> <li>short term use:         increased appetite,         weight gain, mood         change, easy bruising</li> <li>long term use: increase         blood pressure, round         face, osteoporosis</li> </ul>	<ul> <li>Used short term for severe worsening of breathing</li> <li>Take once a day with morning meals</li> <li>If taken longer than 2 weeks, dose may be "tapered" (decrease the amount slowly rather than stop suddenly)</li> </ul>

This material is designed for information purposes only. It should not be used in place of medical advice, instruction and/or treatment.

If you have specific questions, please consult your doctor or appropriate healthcare professional.

<sup>\*</sup>MDI + Meter Dosed Inhaler

<sup>+</sup>Included are most common side effects, may not be complete list.