Calgary COPD & Asthma Program

Medications for Asthma and COPD - Health Care Provider Reference



Category	Trade Name(Generic)	Asthma	COPD	What it Does	Side Effects⁺	Comments
Short Acting Beta Agonist (SABA)	Apo-Salvent Salbutamol Bricanyl [®] (terbutaline) Ventolin [®] (salbutamol) Airomir [®]	X		 rapidly relaxes muscles in the airways works within minutes when needed used before activity to prevent attacks 	 tremor fast heart rate nervousness headache weakness/ dizziness sweating 	 use for sudden shortness of breath use 2 puffs as needed may have to use regularly during flare-ups (Q4H or PRN) use a spacer with all Metered Dose Inhalers (MDIs)
Inhaled Corticosteroid (ICS)	Flovent [®] (fluticasone propionate)	Х	Х	 regular use decreases inflammation of the airways long-term improvement – works over several days CTS guidelines suggests for use in COPD as part of triple therapy for moderate/severe COPD experiencing exacerbations ≥ 1 per year 	hoarse voicesore throatoral thrush	 rinse, gargle, spit, after each use Alvesco[®] can be used OD
	Pulmicort [®] (budesonide)	Х	X			
	Qvar [®] (beclomethasone dipropionate) Alvesco [®] (ciclesonide) Asmanex [®] (mometasone)	X	Χ			use a spacer with all MDIs
		Х	Х			carries a slight increased risk of pneumonia in COPD
		х				
Inhaled Corticosteroid + Long Acting Beta Agonist (ICS + LABA)	Advair [®] (Flovent [®] + Serevent [®])		Х	 decreases inflammation of the airways and relaxes muscles in the airways when taken regularly helps prevent flares of COPD for moderate/severe disease 	 hoarse voice sore throat oral thrush headache tremor fast heart rate muscle cramps 	 rinse, gargle, spit, after each use use a spacer with all MDIs Symbicort[®] may be prescribed as reliever (SMART therapy – asthma. Max. 8 puffs/day) Breo[®] lasts 24 hrs (OD) carries a slight increased risk of pneumonia in COPD CTS guidelines indicate use for mod/severe COPD and ≥1 exacerbation per year* not indicated for mild COPD/Not a First Line Therapy*
	Symbicort [®] (Pulmicort [®] +Oxeze [®])		Х			
	Zenhale [®] (Mometasone+Oxeze ^{®)} Breo [®] (fluticasone furoate+vilanterol)	Х				
		X	X			
Long Acting Beta Agonist (LABA)	Serevent [®] (salmeterol) Oxeze [®] (formoterol) Onbrez [®] (indacaterol maleate)		X	relaxes muscles in the airwaysvaried onset	 headache tremor fast heart rate muscle cramps 	 used as add on therapy <u>with</u> inhaled steroids in asthma Oxeze[®] may be prescribed as reliever for COPD only Onbrez[®] lasts 24 hrs (OD)
Short Acting Muscarinic Antagonist (SAMA)	Atrovent [®] (ipratropium bromide) Combivent [®] (ipratropium bromide + salbutamol)		x	 relaxes muscles in the airways takes about 30 minutes to work lasts 6 hours 	 bad taste or dry mouth tremor (hands shake) headache blurred vision stuffy nose trouble urinating 	 rinse with water if bothered by dry throat or bad taste i do not get in eyes used in the ER for asthma pathway do not use if taking LAMA or dual bronchodialator use a spacer with all MDIs

Calgary COPD & Asthma Program

Medications for Asthma and COPD - Health Care Provider Reference



Category	Trade Name (Generic)	Asthma	COPD	What it Does	Side Effects ⁺	Comments
Long Acting Muscarinic Antagonist (LAMA)	Tudorza [®] (aclidinium bromide) Seebri [®] (glycopyrronium bromide) Spriva [®] (tiotropium bromide)	X	X X X	 relaxes muscles in the airways 	 (May not be complete list) dry mouth constipation trouble urinating headache 	 do not get in eyes If trouble urinating, stop medication and see Doctor Tudorza[®] taken Q12H Spiriva Respimat dosing is 2 inhalations OD
Dual Bronchodilator (LAMA/LABA)	Anoro [®] (umeclidinium+vilanterol) Duaklir [®] (formoteril+aclidinium) Inspiolto [®] (tiotropium+olodaterol) Ultibro [®] (indacaterol+glycopyrronium)		x	 relaxes muscles in the airways lasts 24 hours 	 dry mouth/throat constipation trouble urinating headache tremor fast heart rate muscle cramps 	 once daily use Duaklir taken Q12H do not get in eyes
Oral Corticosteroid	Prednisone	Х		 decreases inflammation of the airways works within hours usually taken for 5-10 days 	 short term: increased appetite, weight gain, mood change, easy bruising long term: increase blood pressure, round face, osteoporosis 	 used short term for severe worsening of breathing take once a day with morning meals if taken longer than 2 weeks, doses are usually "tapered" (decrease the amount slowly rather than stop suddenly).
Leukotriene Antagonist	Singulair[®] (montelukast)	X		 prevents inflammation of the airways 	headachestomach upsetskin rash	 use once per day, in evening
Phosphodiesterase 4 Inhibitor (PDE-4 Inhibitor)	Daxas [®] (roflumilast)		x	 add on therapy to bronchodilator treatment for mod/severe COPD associated with Chronic Bronchitis (cough and sputum) and with frequent exacerbations 	 diarrhea weight decrease nausea headache abdominal pain 	 once daily oral capsule <u>should not</u> be used on its own, or as a rescue medication not recommended for patients with history of depression with suicidal behaviours patient's weight should be monitored regularly not recommended with mod/severe liver impairment
Nasal Corticosteroid	Avamys [®] (fluticasone furoate) Flonase [®] (fluticasone propionate) Nasacort [®] (triamcinolone) Nasonex [®] (mometasone) Rhinocort [®] (budesonide) Omnaris [®] (ciclesonide) beclomethasone)	K	 decreases inflammation in the nose 	 nasal irritation bleeding/ crusting or dry nose 	 use regularly when allergies are bothersome may use as needed throughout the year

This material is designed for information purposes only. It should not be used in place of medical advice, instruction and/or treatment.

*Included are most common side effects, may not be complete list. Information changes frequently – Refer to Canadian Thoracic Society (CTS) http://www.respiratoryguidelines.ca/