

50 or over?

Make your move and ask about

PREVNAR 13

**Don't let the risk of
PNEUMOCOCCAL
PNEUMONIA
ruin your
groove.**



- It can be a serious disease
- Your risk increases after 50

Prevnar® 13 will only protect against diseases caused by the types of *Streptococcus pneumoniae* bacteria covered by the vaccine and it does not reduce complications of pneumococcal disease, such as hospitalization.

Prevnar.ca



Prevnar 13

Your risk increases **with age**

50+ because your immune system naturally weakens as you get older

Your risk also increases **if you are**

18+ with a RISK FACTOR

RISK FACTORS

Medical conditions

- Chronic lung disease (including **asthma** and COPD)
- Chronic heart disease
- Chronic liver disease
- Involving immunosuppression (e.g., transplants, HIV)
- Diabetes
- Cancer
- Certain neurological conditions

Lifestyle factors

- Smoking
 - Alcoholism
 - Living in long-term care facilities
-

Pevnar 13 should not be used if you are allergic (hypersensitive) to the active substances, to any other ingredients in the vaccine, or to any other vaccine that contains diphtheria toxoid.

Take special care with Pevnar 13:

- If you have any present or past medical problems after any dose of Pevnar 13
- If you are sick with a high fever
- If you have any bleeding problems

As with any vaccine, Pevnar 13 will not protect 100% of those who receive the vaccine.