

An Atlas of Lumps and Bumps, Part 49: Solitary Cutaneous Mastocytoma


 consultant360.com/photo-essay/atlas-lumps-and-bumps-part-49-solitary-cutaneous-mastocytoma

Photo Essay

Volume 65 - Issue 3 - March 2025

Solitary Cutaneous Mastocytoma

Mastocytosis is a rare heterogenous group of disorders characterized by a clonal proliferation and accumulation of mast cells in one or more organs, mostly skin and bone marrow and, less frequently, gastrointestinal tract, liver, spleen, and lymph nodes.¹⁻³ Depending on the sites of organ involvement, two main forms of mastocytosis are recognized, namely, cutaneous mastocytosis (when mast cell accumulation is limited to the skin) and systemic mastocytosis (when mast cell accumulation involves extracutaneous tissues particularly the bone marrow, often combined with skin involvement).⁴ Cutaneous mastocytosis can be subclassified into one of the three subtypes, namely, solitary cutaneous mastocytoma, maculopapular cutaneous mastocytosis (also known as urticaria pigmentosa), and diffuse cutaneous mastocytosis.⁵

Somatic activating mutations in the *c-KIT* proto-oncogene (most commonly D816V) that encodes the mast cell growth receptor KIT (also called CD117), a transmembrane protein that binds to stem cell factor and promotes cell division, have been shown to result in abnormal proliferation of mast cells and melanocytes.^{1,2,6-9} A proliferation of melanocytes accounts for the hyperpigmented lesions seen in cutaneous mastocytosis.^{8,10,11} Disease manifestations such as pruritus, flushing, blistering, wheals, vomiting, abdominal pain, hypotension, and dyspnea are caused by spontaneous or induced activation of mast cells with resulting release of vasoactive mediators.^{12,13}

The prevalence of mastocytosis has been estimated to be 1 in 10,000; 80% of these patients have cutaneous mastocytosis.^{1,13} Solitary cutaneous mastocytoma accounts for 10 to 15% of children with cutaneous mastocytosis.^{1,14,15} There is no gender or race predilection.¹⁶

Solitary cutaneous mastocytoma most often presents at birth or develops within the first few months of life.^{4,13,14} Typically, solitary cutaneous mastocytoma presents with an indurated, nontender, erythematous, yellow (**Figure 1**), orange (**Figure 2**), or brown (**Figure 3**), oval or round macule, papule, plaque or nodule.^{1,17,18} The lesion is usually 1 to 5 cm in diameter and often has a peau

d'orange (pebbly, orange peel-like) appearance and a leathery or rubbery consistency.^{1,6,17} The lesional margins can be sharp or indistinct.⁵



Fig. 1. The prevalence of mastocytosis has been estimated to be 1 in 10,000. Occasionally, more than one mastocytoma (up to a maximum of three) can be found.



Fig. 2. Solitary cutaneous mastocytoma accounts for 10 to 15% of children with cutaneous mastocytosis.



Fig. 3. Typically, solitary cutaneous mastocytoma presents with an indurated, nontender, erythematous, yellow, orange, or brown, oval or round macule, papule, plaque or nodule.

Occasionally, more than one mastocytoma (up to a maximum of three) can be found.^{4,2,13} Pruritus is variable; the lesion can be asymptomatic or moderately pruritic.^{1,12,17} The lesion usually increases in size (but not in number) for several months and then grows in proportion to the size of the child for a variable period of time before regressing over time.^{1,19} Sites of predilection include the trunk, extremities, axillae, groins, followed by the head and neck.^{1,19} The palms and soles are usually spared.⁶ The lesion may spontaneously urticate and/or, more commonly, urticate when rubbed or stroked.^{1,19,20}

Upon rubbing or stroking, the lesion becomes pruritic, erythematous, or edematous; this reaction is referred to as Darier sign (**Figure 4**).^{1,19,21} The Darier sign is caused by degranulation of mast cells with subsequent release of mast cell mediators induced by physical stimulation. Although the Darier sign is pathognomonic,^{16,21} it is only seen in 50% of patients with solitary cutaneous mastocytoma (**Figure 5**).^{1,22,23}

As such, the absence of a Darier sign does not exclude the diagnosis of a solitary cutaneous mastocytoma.²

Some of the lesions may blister with irritation.^{1,24}

Although systemic symptoms such as flushing, dyspnea, hypotension, nausea, vomiting, abdominal pain, diarrhea, and headache are much more commonly seen in patients with systemic mastocytosis, they can also occur in patients with solitary cutaneous mastocytoma.^{1,24} Therefore, large or extensive lesion of solitary cutaneous mastocytoma should not be vigorously rubbed and the Darier sign should not be elicited in case of a large mastocytoma as such a maneuver can precipitate severe symptoms.^{1,25}

Organomegaly and lymphadenopathy are characteristically absent.^{1,25}



Fig. 4. Upon rubbing or stroking, the lesion becomes pruritic, erythematous, or edematous, which is known as the Darier sign.



Fig. 5. The Darier sign is only seen in 50% of patients with solitary cutaneous mastocytoma.

The diagnosis is mainly clinical, based on the morphology of the lesion, the presence of a positive Darier sign, and absence of systemic involvement.^{1,6}

Dermoscopic features of a solitary cutaneous mastocytoma include light-brown blots, reticular vascular pattern, yellow-orange blots (attributable to dense papillary and reticular dermal mast cell infiltrate), and a

pigment network (attributable to dermal mast cell infiltrate and accumulation of melanin in basal keratinocytes).^{20,23} When Darier sign is present, dermoscopy typically shows a decrease in yellowish hue and pigment network intensity along with the appearance of peripheral erythema.²³ A skin biopsy is not essential but can be helpful if the diagnosis is in doubt.

Generally, the prognosis is good. Most children whose onset of disease is before two years of age tend to have spontaneous regression of skin lesions by 6 years of age and resolution of the disease by 10 years of age.^{1,13,15,16}

AUTHORS:

Alexander K.C. Leung, MD^{1,2}, Benjamin Barankin, MD³, Joseph M. Lam, MD⁴, Kin Fon Leong, MD⁵

AFFILIATIONS:

¹Clinical Professor of Pediatrics, the University of Calgary, Calgary, Alberta, Canada

²Pediatric Consultant, the Alberta Children's Hospital, Calgary, Alberta, Canada

³Dermatologist, Medical Director and Founder, the Toronto Dermatology Centre, Toronto, Ontario, Canada

⁴Associate Clinical Professor of Pediatrics, Dermatology and Skin Sciences, the University of British Columbia,

Vancouver, British Columbia, Canada.

⁵Pediatric Dermatologist, the Pediatric Institute, Kuala Lumpur General Hospital, Kuala Lumpur, Malaysia

CITATION:

Leung AKC, Barankin B, Lam JM, Leong KF. An Atlas of Lumps and Bumps, Part 49: Solitary Cutaneous Mastocytoma. *Consultant*. 2025;65(3).
doi: 10.25270/con.2025.03.000001

CORRESPONDENCE:

Alexander K. C. Leung, MD, #200, 233 16th Ave NW, Calgary, AB T2M 0H5, Canada (aleung@ucalgary.ca)

EDITOR'S NOTE:

This article is part of a series describing and differentiating dermatologic lumps and bumps. To access previously published articles in the series, visit: <https://www.consultant360.com/resource-center/atlas-lumps-and-bumps>.

References

1. Leung AKC, Lam JM, Leong KF. Childhood solitary cutaneous mastocytoma: Clinical manifestations, diagnosis, evaluation, and management. *Curr Pediatr Rev*. 2019;15(1):42-46.
doi:10.2174/1573396315666181120163952.

2. Brockow K. Urticaria pigmentosa. *Immunol Allergy Clin North Am*. 2004;24(2):287-316.
doi:10.1016/j.iac.2004.01.002.
3. Tzankov A, Duncavage E, Craig FE, et al. Mastocytosis. *Am J Clin Pathol*. 2021;155(2):239-266. doi:10.1093/ajcp/aqaa183.
4. Matito A, Azaña JM, Torrelo A, Alvarez-Twose I. Cutaneous mastocytosis in adults and children: New classification and prognostic factors. *Immunol Allergy Clin North Am*. 2018;38(3):351-363.
doi:10.1016/j.iac.2018.04.001.
5. Hartmann K, Escribano L, Grattan C, et al. Cutaneous manifestations in patients with mastocytosis: consensus report of the European Competence Network on Mastocytosis; the American Academy of Allergy, Asthma & Immunology; and the European Academy of Allergology and Clinical Immunology. *J Allergy Clin Immunol*. 2016;137(1):35-45.
doi:10.1016/j.jaci.2015.08.034.
6. Azana JM, Torrelo A, Matito A. Update on mastocytosis (part 1): pathophysiology, clinical features, and diagnosis. *Actas Dermosifiliogr*. 2016;107(1):5-14. doi:10.1016/j.ad.2015.09.007.
7. Castells M, Akin C. Mastocytosis (cutaneous and systemic): epidemiology, pathogenesis, and clinical manifestations. In: Post TW, ed. *UpToDate*. Waltham, MA: UpToDate; 2021. Accessed March 12, 2021.

8. Cowart D, Cowart K, Hougeir FG. A 39-year-old woman with spots covering her body. *BMJ Case Rep.* 2013;2013:bcr2013201434. doi:10.1136/bcr-2013-201434.
9. Wiechers T, Rabenhorst A, Schick T, et al. Large maculopapular cutaneous lesions are associated with favorable outcome in childhood-onset mastocytosis. *J Allergy Clin Immunol.* 2015;136(6):1581-1590.e3. doi:10.1016/j.jaci.2015.05.034.
10. Miller MD, Nery NS, Gripp AC, Maceira JP, Nascimento GM. Dermatoscopic findings of urticaria pigmentosa. *An Bras Dermatol.* 2013;88(6):986-988. PMID: 24474113.
11. Peters F, Fiebig B, Lundberg P, et al. Detection of the germline KIT S476I mutation in a kindred with familial mastocytosis associated with gastrointestinal stromal tumors. *J Allergy Clin Immunol Pract.* 2021:S2213-2198(21)00004-0. doi:10.1016/j.jaip.2020.12.049.
12. Castells M, Akin C. Mastocytosis (cutaneous and systemic): Evaluation and diagnosis in children. In: Post TW, ed. *UpToDate*. Waltham, MA: UpToDate; 2021. Accessed March 12, 2021.
13. Di Raimondo C, Del Duca E, Silvaggio D, et al. Cutaneous mastocytosis: A dermatological perspective. *Australas J Dermatol.* 2021;62(1):e1-e7. doi:10.1111/ajd.13443.

14. Ha NH, Lee YJ, Park MC, et al. Solitary mastocytoma presenting at birth. *Arch Craniofac Surg*. 2018;19(2):127-130. doi:10.7181/acfs.2018.01508.
15. Hussain SH. Pediatric mastocytosis. *Curr Opin Pediatr*. 2020;32(4):531-538. doi:10.1097/MOP.0000000000000922.
16. Ahmed A, Jan A. Mastocytoma. In: *StatPearls [Internet]*. Treasure Island, FL: StatPearls Publishing; 2021. PMID: 30855840.
17. Gopal D, Puri P, Singh A, Ramesh V. Asymptomatic solitary cutaneous mastocytoma: a rare presentation. *Indian J Dermatol*. 2014;59(6):634. doi:10.4103/0019-5154.143588.
18. Exposito-Serrano V, Agut-Busquet E, Leal Canosa L, et al. Pleomorphic mastocytoma in an adult. *J Cutan Pathol*. 2018;45(2):176-179. doi:10.1111/cup.13080.
19. Gori A, Torneria C, Kelly VM, Zlotoff BJ, Contreras ME. Visual diagnosis: two infants who have skin lesions that react to minor trauma. *Pediatr Rev*. 2009;30(7):280-283. doi:10.1542/pir.30-7-280.
20. Kumar A, Kansal NK, Anuragi RP. Cutaneous mastocytosis: clinical, dermoscopic and pathological features. *BMJ Case Rep*. 2020;13(12):e240169. doi:10.1136/bcr-2020-240169.

21. de Giorgi V, Fabroni C, Alfaioli B, et al. Solitary mastocytoma: tooth eruption as triggering factor. *Int J Dermatol*. 2008;47(12):1274-1277. doi:10.1111/j.1365-4632.2008.03841.x.
22. Conrad H, Gausche-Hill M, Burbulys D. A 6-month-old with total body flushing and a macular-papular lesion. *Pediatr Emerg Care*. 2007;23(5):321-323. doi:10.1097/01.pec.0000270165.85862.12.
23. Mukherjee S, Neema S, Sinha P, et al. Dermoscopic demonstration of Darier sign. *Dermatol Pract Concept*. 2021;11(1):e2021114. doi:10.5826/dpc.1101a114.
24. Bulat V, Mihić LL, Situm M, et al. Most common clinical presentations of cutaneous mastocytosis. *Acta Clin Croat*. 2009;48(1):59-64. PMID: 19623875.
25. Klaiber N, Kumar S, Irani AM. Mastocytosis in children. *Curr Allergy Asthma Rep*. 2017;17(11):80. doi:10.1007/s11882-017-0748-4.