

Frequently Asked Questions (FAQs)

Inpatient/PICU – Children’s Hospital of Eastern Ontario

Q: What is this study all about?

A: The PRoMPT BOLUS study will determine if resuscitation with balanced IV fluids (Lactated Ringers) results in improved outcomes in children aged 2 months < 18 years presenting to the Emergency Department with sepsis.

Q: Will children who become septic on the ward/in the PICU be eligible?

A: No. This study is enrolling exclusively in the ED. This is to avoid any confounding from fluids previously administered on the ward, in the PICU, or during transport.

Q: How will I know if my patient is enrolled in the study?

A: The ED Clinical staff will inform the admitting team about enrollment during handover. We are using multiple colour-coded visual cues (**Green = Normal Saline**, **Pink = Lactated Ringers**) to let you know about the study:

- 1) Patient Wrist Bands
- 2) IV hang tags

In the MAR, treating team members will also see the “PRoMPT Study” beside the name of the IV fluid.

Q: Will I know which fluids my patient is allocated to?

A: Yes. This is an open-label study (i.e. not blinded).

Multiple colour-coded visual cues (**Green = Normal Saline**, **Pink = Lactated Ringers**) will remind you what fluid the participant has been allocated to.

Q: How much IV fluid should I order?

A: The amount, timing, and rate of fluid administered is at the discretion of the health care provider. The study fluids includes both bolus and maintenance fluids and includes medication infusions as much as possible (see below).

Q: Are there special study-specific fluid bags?

A: No. You will use the fluids that are available as part of the routine hospital stock.

Q: Can I use a different fluid than the one my patient is allocated to?

A: Adherence to the allocated fluid is very important to maintain the quality of the evidence derived from the study. However, if there is a *definitive clinical indication* for using an alternate fluid from the one allocated, please manage the patient based on that clinical requirement (note: should be rare/exceptional). An example would be using saline for clinically relevant hyponatremia.

Q: Can I add electrolytes, dextrose, etc?

A: Yes - Some patients will require added electrolytes, dextrose etc.

Q: What about mixing medication infusions?

A: Please follow routine pharmacy directives for mixing medications, regardless of what study fluid your patient has been allocated to.

Q: Are there any compatibility issues?

A: There are several well-known compatibility issues for Lactated Ringers (LR). LR cannot be provided through the same IV line at the same time as ceftriaxone or blood products. Options

include the use of a second IV line, or a very careful IV flush prior to- and after- the use of these medications. Please review and follow all CHEO pharmacy administration guidelines for all medication administration and compatibility concerns.

Q: When does the study intervention end?

A: The study intervention period continues until 11:59pm the day following randomization.

The allocated fluid should be provided until that point. Following the intervention period, you can continue the allocated fluid should you choose, or switch to any fluid of your choice.

Note: the date/time of randomization will be written on the IV Hang Tags

Q: What about other interventions like antibiotics, vasoactive medications etc?

A: This study only controls the type of fluid administered to the child. ALL other interventions are to be done according to best practice at the discretion of the managing health care provider.

Q: Are there any reasons I should stop the study fluids early?

A: There are several rare but concerning clinical situations in which a child should be stopped from the study. These include:

- 1) Significant electrolyte abnormalities such as hyperkalemia and hypercalcemia, **confirmed on repeat testing**
- 2) Significant end organ damage such as severe hepatic or renal impairment,
- 3) Hypersensitivity to the allocated fluid

Formal definitions can be found on the PRoMPT BOLUS website. See QR code below.

Q: What should I do if a child meets Stopping Criteria?

A: If a child meets one of the Stopping Criteria, they are no longer required to receive the allocated study fluid. Manage the child at your discretion, and when possible, notify the study team.

Q: What about consent?

A: Due to the emergent nature of sepsis, and the need for immediate administration of fluid management, it is not feasible to obtain true informed consent at the time of fluid initiation in the ED. This study has been approved by our local REB for deferred consent. A Caregiver Information Sheet was provided to the family in the ED. A member of the research team will approach the family at the most optimal time to further discuss the study and obtain deferred consent.

Q: HELP!!! I have questions about the patient in front of me...what do I do?

A: Please refer to the following resources for more help:

- 1) ED Study Information Sheet
- 2) Contact a study team member:
 - Dr. Waleed Alqurashi (walqurashi@cheo.on.ca)
 - Dr. Fuad Alnaji (fnaji@cheo.on.ca)
 - ED Study Coordinator: Cheryl Kreviazuk (ckreviazuk@cheo.on.ca)
 - PICU Study Coordinator: Katie O'Hearn (kohearn@cheo.on.ca)

