Inpatient/PICU Team Instructions



| Review admitted patients to confirm if they are a PRoMPT BOLUS participant. Identify fluid allocation on |
|--|
| patient bracelet and/or IV hang tags. |
| Decimands and transfer and assistance of an additional and additional additional and additional additiona |

- Documents and treatment assignment are colour coded
 - Yellow = 0.9% Normal Saline (NS)
 - Green = Balanced Fluid (BF: Lactated Ringers or PlasmaLyte)
- Keep coloured IV hang tag and coloured patient bracelet with the patient
- ☐ Place order for **ALL boluses and maintenance fluid** provided in ED:
 - Order the fluid TYPE (NS or BF) specified by randomization information
 - Note: Hypotonic fluids (i.e. 0.45NS) are not considered appropriate for bolus or maintenance fluids in septic children and should be avoided
 - When and how much fluid should be decided based on clinical need by the clinical care team
 - Obtain fluids from clinical ED inventory
 - Additives (i.e. Dextrose, KCI) can be used as needed
 - Document <u>ALL IV and IO fluid given</u> to the patient during this time
- □ During the patient transfer of care/handover verbally inform the receiving team that the child is participating in the PRoMPT BOLUS study
 - Inpatient/PICU teams should continue to order the allocated study fluid TYPE (NS or BF)
 through 23:59 on the calendar day following randomization
 - Keep coloured IV hang tag and coloured patient bracelet with the patient
- ☐ If patient meets Stopping Criteria, manage the child at your discretion, and:
 - Notify the study team that the child had been enrolled but met stopping criteria
- ☐ Any questions, please contact
 - Dr. Gary Joubert at Gary.Joubert@lhsc.on.ca or (519) 851-7989
 - Dr. Gurinder Sangha at Gurinder.Sangha@lhsc.on.ca or #15556
 - Study team pager (Mon-Fri, 08:00-16:00): #17561 / Patrick.Siedlecki@lhsc.on.ca



