## **ED Team Instructions**

## BC Children's Hospital



□ Complete study Eligibility Form in full:

- Sign and Date form as soon as randomization occurs
- Affix randomization & patient label to eligibility form
- □ Identify Fluid Allocation:
  - Fluid outlined in this package is the fluid to be administered (colour coded)
    - 0.9% Normal Saline (NS yellow)
    - Balanced Fluids (BF green) Lactated Ringers OR PlasmaLyte (clinician preference)

□ Place order in CST – For <u>ALL boluses and maintenance fluid</u> provided in the ED:

- Order the fluid type (NS or BF) specified by the randomization information
- <u>Note:</u> Hypotonic fluids (i.e. 0.45NS) are not considered appropriate for bolus or maintenance fluids in septic children and should be avoided
- When and how much fluid should be decided based on clinical need by the clinical care team.
- Obtain fluids from clinical ED inventory
- Similarly for maintenance fluids, please use D5 Normal or D5 Ringers as per their randomization.
- Physician to place a "Research Communication to Nursing" on CST stating: "Patient enrolled in PROMPT BOLUS study and randomized to [insert Normal Saline or Balanced Fluids]"
- □ Write **<u>date and time of fluid start</u>** on IV hang tag and door signs where indicated:
  - Hang IV Hang Tag on pole
- $\hfill\square$  Place the patient bracelet on the patient wrist
- □ Provide Caregiver with the Caregiver Information Sheet:
  - This study is approved to be conducted under a deferred consent model
  - The patient/caregiver will not consent before randomization/fluid administration occur
  - A member of the study team will approach the family when it is clinically appropriate
- □ Communicate with admitting team:
  - Verbally inform the admitting team that the child is participating in the PRoMPT BOLUS Study and which fluids they have been allocated to
  - Provide the admitting team with Inpatient Instruction Sheet
  - Inpatient teams should continue to order study fluid TYPE (NS or BF) through 23:59 on the calendar day following randomization.
- □ If patient is discharged from ED discontinue fluids at discharge and place any forms in the physical chartlet

□ Place Inpatient Instructions with the patient physical chartlet <u>BEFORE</u> transfer or admission



□ More information available on study website - use QR Code: